

LEUCOVORIN CALCIUM - leucovorin calcium tablet

Roxane Laboratories, Inc


RX ONLY

DESCRIPTION

Leucovorin Calcium Tablets USP contain either 5 mg, 10 mg, 15 mg or 25 mg leucovorin as the calcium salt of *N*-[4-[[[(2-amino-5-formyl-1,4,5,6,7,8-hexahydro-4-oxo-6-pteridiny]methyl] amino]benzoyl]-*L*-glutamic acid. This is equivalent to either 5.4 mg, 10.8 mg, 16.21 mg or 27.01 mg of anhydrous leucovorin calcium, respectively. In addition, each tablet contains the following *inactive ingredients*: microcrystalline cellulose, corn starch, croscarmellose sodium, povidone, colloidal silicon dioxide, magnesium stearate, D&C yellow #10 (15 mg and 25 mg).

Leucovorin is a water soluble form of reduced folate in the folate group; it is useful as an antidote to drugs which act as folic acid antagonists. These tablets are intended for oral administration only.

The structural formula of leucovorin calcium is:



C₂₀H₂₁CaN₇O₇ M.W. 511.51

CLINICAL PHARMACOLOGY

Leucovorin is a racemic mixture of the diastereoisomers of the 5-formyl derivative of tetrahydrofolic acid. The biologically active compound of the mixture is the (-)-*L*-isomer, known as *Citrovorum factor*, or (-)-folinic acid. Leucovorin does *not* require reduction by the enzyme dihydrofolate reductase in order to participate in reactions utilizing folates as a source of “one-carbon” moieties. Following oral administration, leucovorin is rapidly absorbed and enters the general body pool of reduced folates. The increase in plasma and serum folate activity (determined microbiologically with *Lactobacillus casei*) seen after oral administration of leucovorin is predominantly due to 5-methyltetrahydrofolate.

Twenty normal men were given a single, oral 15 mg dose (7.5 mg/m²) of leucovorin calcium and serum folate concentrations were assayed with *L. casei*. Mean values observed (\pm one standard error) were:

- Time to peak serum folate concentration: 1.72 \pm 0.08 hours,
- Peak serum folate concentration achieved: 268 \pm 18 ng/mL,
- Serum folate half-disappearance time: 3.5 hours.

Oral tablets yielded areas under the serum folate concentration-time curves (AUCs) that were 12% greater than equal amounts of leucovorin given intramuscularly and equal to the same amounts given intravenously.

Oral absorption of leucovorin is saturable at doses above 25 mg. The apparent bioavailability of leucovorin was 97% for 25 mg, 75% for 50 mg and 37% for 100 mg.

INDICATIONS AND USAGE

Leucovorin Calcium Tablets USP are indicated to diminish the toxicity and counteract the effects of impaired methotrexate elimination and of inadvertent overdoses of folic acid antagonists.

CONTRAINDICATIONS

Leucovorin is improper therapy for pernicious anemia and other megaloblastic anemias secondary to the lack of vitamin B₁₂. A hematologic remission may occur while neurologic manifestations continue to progress.

WARNINGS

In the treatment of accidental overdosage of folic acid antagonists, leucovorin should be administered as promptly as possible. As the time interval between antifolate administration (e.g., methotrexate) and leucovorin rescue increases, leucovorin's effectiveness in counteracting hematologic toxicity decreases.

Monitoring of the serum methotrexate concentration is essential in determining the optimal dose and duration of treatment with leucovorin.

Delayed methotrexate excretion may be caused by a third space fluid accumulation (i.e., ascites, pleural effusion), renal insufficiency, or inadequate hydration. Under such circumstances, higher doses of leucovorin or prolonged administration may be indicated. Doses higher than those recommended for oral use must be given intravenously.

Leucovorin may enhance the toxicity of fluorouracil. Deaths from severe enterocolitis, diarrhea, and dehydration have been reported in elderly patients receiving weekly leucovorin and fluorouracil.¹ Concomitant granulocytopenia and fever were present in some but not all of the patients.

The concomitant use of leucovorin with trimethoprim-sulfamethoxazole for the acute treatment of *Pneumocystis carinii* pneumonia in patients with HIV infection was associated with increased rates of treatment failure and mortality in a placebo-controlled study.

PRECAUTIONS

General

Parenteral administration is preferable to oral dosing if there is a possibility that the patient may vomit or not absorb the leucovorin. Leucovorin has no effect on other established toxicities of methotrexate such as the nephrotoxicity resulting from drug and/or metabolite precipitation in the kidney.

Drug Interactions

Folic acid in large amounts may counteract the antiepileptic effect of phenobarbital, phenytoin and primidone, and increase the frequency of seizures in susceptible children.

Preliminary animal and human studies have shown that small quantities of systemically administered leucovorin enter the CSF primarily as 5-methyltetrahydrofolate and, in humans, remain 1 to 3 orders of magnitude lower than the usual methotrexate concentrations following intrathecal administration. However, high doses of leucovorin may reduce the efficacy of intrathecally administered methotrexate.

Leucovorin may enhance the toxicity of fluorouracil (see WARNINGS).

Pregnancy

Teratogenic Effects

Pregnancy Category C

Animal reproduction studies have not been conducted with leucovorin. It is also not known whether leucovorin can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Leucovorin should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when leucovorin is administered to a nursing mother.

Pediatric Use

See Drug Interactions subsection.

ADVERSE REACTIONS

Allergic sensitization, including anaphylactoid reactions and urticaria, has been reported following the administration of both oral and parenteral leucovorin.

OVERDOSAGE

Excessive amounts of leucovorin may nullify the chemotherapeutic effect of folic acid antagonists.

DOSAGE AND ADMINISTRATION

Leucovorin calcium tablets are intended for oral administration. Because absorption is saturable, oral administration of doses greater than 25 mg is not recommended.

Impaired Methotrexate Elimination or Inadvertent Overdosage

Leucovorin rescue should begin as soon as possible after an inadvertent overdosage and within 24 hours of methotrexate administration when there is delayed excretion (see WARNINGS). Leucovorin 15 mg (10 mg/m²) should be administered IM, IV, or PO every 6 hours until the serum methotrexate level is less than 10⁻⁸ M. In the presence of gastrointestinal toxicity, nausea, or vomiting, leucovorin should be administered parenterally.

Serum creatinine and methotrexate levels should be determined at 24-hour intervals. If the 24-hour serum creatinine has increased 50% over baseline or if the 24-hour methotrexate level is greater than 5 x 10⁻⁶ M or the 48-hour level is greater than 9 x 10⁻⁷ M, the dose of leucovorin should be increased to 150 mg (100 mg/m²) IV every 3 hours until the methotrexate level is less than 10⁻⁸ M. Doses greater than 25 mg should be given parenterally (see CLINICAL PHARMACOLOGY).

Hydration (3 L/d) and urinary alkalization with sodium bicarbonate should be employed concomitantly. The bicarbonate dose should be adjusted to maintain the urine pH at 7.0 or greater.

The recommended dose of leucovorin to counteract hematologic toxicity from folic acid antagonists with less affinity for mammalian dihydrofolate reductase than methotrexate (i.e., trimethoprim, pyrimethamine) is substantially less, and 5 to 15 mg of leucovorin per day has been recommended by some investigators.

Patients who experience delayed early methotrexate elimination are likely to develop reversible non-oliguric renal failure. In addition to appropriate leucovorin therapy, these patients require continuing hydration and urinary alkalization, and close monitoring of fluid and electrolyte status, until the serum methotrexate level has fallen below 0.05 micromolar and the renal failure has resolved.

Some patients will have abnormalities in methotrexate elimination or renal function following methotrexate administration, which are significant but less severe. These abnormalities may or may not be associated with significant clinical toxicity. If significant clinical toxicity is observed, leucovorin rescue should be extended for an additional 24 hours (total 14 doses over 84 hours) in subsequent courses of therapy. The possibility that the patient is taking other medications which interact with methotrexate (e.g., medications which may interfere with methotrexate elimination or binding to serum albumin) should always be reconsidered when laboratory abnormalities or clinical toxicities are observed.

HOW SUPPLIED

Leucovorin Calcium Tablets USP are available for oral administration as:

5 mg white, scored tablets (Identified 54 293).

NDC 0054-8496-19: Unit dose, 10 tablets per strip, 5 strips per shelf pack.

NDC 0054-4496-13: Bottles of 30 tablets.

NDC 0054-4496-25: Bottles of 100 tablets.

10 mg white, scored tablets (Identified 54 942).

NDC 0054-4497-05: Bottles of 12 tablets.

NDC 0054-4497-10: Bottles of 24 tablets.

15 mg yellow, scored tablets (Identified 54 650).

NDC 0054-4498-10: Bottles of 24 tablets.

25 mg yellow, scored tablets (Identified 54 013).

NDC 0054-4499-11: Bottles of 25 tablets.

Store at Controlled Room Temperature 15° to 30°C (59° to 86°F).

Protect From Light and Moisture.

References:

1.Grem JL, Shoemaker DD, Petrelli NJ, Douglas HO. Severe and fatal toxic effects observed in treatment with high- and low-dose leucovorin plus 5-fluorouracil for colorectal carcinoma. Cancer Treat Rep 1987;71:1122.

2.Link MP, Goorin AM, Miser AW et al. The effect of adjuvant chemotherapy on relapse-free survival in patients with osteosarcoma of the extremity. N Engl J Med 1986;314:1600-1606.

4055309//03 Revised December 2007

© RLI, 2007

PACKAGE LABEL - LEUCOVORIN CALCIUM TABLETS USP

5 mg white, scored tablets (Identified 54 293).

NDC 0054-8496-19: Unit dose, 10 tablets per strip, 5 strips per shelf pack.

Rx Only

Roxane Laboratories, Inc.

N 3 0054-8496-19 9

NDC 0054-8496-19 5x10 Unit-Dose Tablets

**LEUCOVORIN
CALCIUM
Tablets USP**

5 mg*

*Each tablet contains leucovorin calcium equivalent to 5 mg leucovorin.

See Package Insert for Complete Prescribing Information.

Store at Controlled Room Temperature 15° to 30°C (59° to 86°F).

Protect From Light and Moisture.

This unit-dose package is not child-resistant. If dispensed for outpatient use, a child-resistant container should be utilized.

R_x only

TABLETS IDENTIFIED 54 293

Roxane Laboratories, Inc., Columbus, Ohio 43216

NDC 0054-8496-19 5x10 Unit-Dose Tablets

**LEUCOVORIN
CALCIUM
Tablets USP**

5 mg*

LOT

EXP.

 **Boehringer Ingelheim**
Roxane Laboratories

4254610//05
© RLI, 2007

NDC 0054-4496-13: Bottles of 30 tablets.

Rx Only

Roxane Laboratories, Inc.

See Package Insert for Complete Prescribing Information.

Store at Controlled Room Temperature 15° to 30°C (59° to 86°F).

Protect From Light and Moisture.

Dispense in a light-resistant container as defined in the USP/NF.

TABLETS IDENTIFIED 54 293

Roxane Laboratories, Inc.
Columbus, Ohio 43216

NDC 0054-4496-13 30 Tablets

**LEUCOVORIN
CALCIUM
Tablets USP**

5 mg*

*Each tablet contains leucovorin calcium equivalent to 5 mg leucovorin.

R_x only

 **Roxane Laboratories**

4178942//03
© RLI, 2007

NDC 0054-4496-25: Bottles of 100 tablets.

Rx Only

Roxane Laboratories, Inc.

See Package Insert for Complete Prescribing Information.

Store at Controlled Room Temperature 15° to 30°C (59° to 86°F).

Protect From Light and Moisture.

Dispense in a light-resistant container as defined in the USP/NF.

TABLETS IDENTIFIED 54 293

Roxane Laboratories, Inc.
Columbus, Ohio 43216

NDC 0054-4496-25 100 Tablets

**LEUCOVORIN
CALCIUM
Tablets USP**

5 mg*

*Each tablet contains leucovorin calcium equivalent to 5 mg leucovorin.

R_x only

 **Roxane Laboratories**

4178942//03
© RLI, 2007

10 mg white, scored tablets (Identified 54 942).

NDC 0054-4497-05: Bottles of 12 tablets.

Rx Only

Roxane Laboratories, Inc.

See Package Insert for Complete Prescribing Information.

Store at Controlled Room Temperature 15° to 30°C (59° to 86°F).

Protect From Light and Moisture.

Dispense in a light-resistant container as defined in the USP/NF.

TABLETS IDENTIFIED 54 942

Roxane Laboratories, Inc.
Columbus, Ohio 43216

NDC 0054-4497-05 12 Tablets

**LEUCOVORIN
CALCIUM
Tablets USP**

10 mg*

*Each tablet contains leucovorin calcium equivalent to 10 mg leucovorin.

R_x only

 **Roxane Laboratories**

4178948//03
© RLI, 2007

NDC 0054-4497-10: Bottles of 24 tablets.

Rx Only

Roxane Laboratories, Inc.



15 mg yellow, scored tablets (Identified 54 650).

NDC 0054-4498-10: Bottles of 24 tablets.

Rx Only

Roxane Laboratories, Inc.



25 mg yellow, scored tablets (Identified 54 013).

NDC 0054-4499-11: Bottles of 25 tablets.

Rx Only

Roxane Laboratories, Inc.

